

Effects of Vertebral Axial Decompression (VAX-D) On Intradiscal Pressure

Gustavo Ramos, M.D., William Marin, M.D.

Journal of Neurosurgery 81:350-353 1994

Departments of Neurosurgery and Radiology, Rio Grande Regional Hospital, McAllen, and
Division of Neurosurgery, Health Sciences Center,
University of Texas, San Antonio, Texas

Abstract

The object of this study was to examine the effect of vertebral axial decompression on pressure in the nucleus pulposus of lumbar discs. Intradiscal pressure measurement was performed by connecting a cannula inserted into the patient's L4-5 disc space to a pressure transducer. The patient was placed in a prone position on a VAX-D therapeutic table and the tensionometer on the table was attached via a pelvic harness. Changes in intradiscal pressure were recorded at resting state and while controlled tension was applied by the equipment to the pelvic harness. Intradiscal pressure demonstrated an inverse relationship to the tension applied. Tension in the upper range was observed to decompress the nucleus pulposus significantly, to below -100 mm Hg.

KEY WORDS - low-back pain, intradiscal pressure, vertebral decompression

Introduction

Surgical procedures utilizing conventional and percutaneous approaches have established the merits of decompression of intravertebral disc spaces in the management of low-back pain syndrome associated with lumbar disc herniation.^{4,12,13,15} Surgery will continue to play an important role in the treatment of patients with low-back pain and sciatica associated with herniated discs and degenerative disc problems. However, for patients who are not candidates for surgery, there is a need to establish a conservative approach that offers an effective means of returning the patient to a functional level of activity.

Considerable controversy exists in regard to the various techniques currently employed. Aside from basic bed rest, there are few noninterventional modalities that have been adopted as standards of therapy. Manipulative techniques for mechanical low-back pain associated with posterior facet syndrome or muscle strain have not been found as useful in the management of herniated or degenerated lumbar discs. Similarly, other modalities including ultrasound treatments, various electrical stimulation techniques, short-wave therapy acupuncture, steroid injections, and the administration of anti-inflammatory agents and muscle relaxants all have a following among some practitioners but fall short of addressing the underlying problems associated with intervertebral disc lesions. All of these treatment methods fail by comparison to surgery, in our opinion, because they have the common problem of not relieving the pain from neurocompression or from the stimuli associated with a prolapsed nucleus pulposus. The only noninterventional method that has been shown to

hold any promise of relieving pressure on vital structures of the lumbar region is that of distraction of the lumbar vertebrae by mechanical forces applied along the axis of the spinal column.^{2,3,5,14}

There has been some investigation into the effects of distracting segments of the spinal column excised from cadavers,^{11,14} as well as radiological studies that provided evidence that the application of certain forms of tension can distract vertebral bodies.^{3,5} On the other hand, there are equally pertinent studies that failed to demonstrate any positive effects from other methods of applying spinal traction.^{1,10} Nachemson and Elfstrom⁶⁻⁹ have studied the effects of movement and posture on intradiscal pressure. Their measurements show pressure changes caused by positioning and posture range between 25 and 275 mm Hg, suggesting that some positions and postures may be inadvisable for patients suffering from lumbar disc lesions. Anderson, et al.¹ and others have shown that certain traction techniques can actually cause an increase in intradiscal pressure, which would be undesirable in the treatment of low-back pain associated with herniated discs and a neurocompression etiology.

A new form of therapy, termed "vertebral axial decompression," has recently been introduced in the physical therapy department of the Rio Grande Regional Hospital. This treatment modality has shown considerable promise in relieving low-back pain associated with herniated discs or degenerative disc disease of the lumbar vertebrae in patients who are not considered candidates for surgery. The purpose of this research project was to investigate the influence of this new treatment modality on intradiscal pressure in the lumbar spine of patients receiving this form of therapy.

Clinical Material and Methods

Five cases were selected from among individuals who were referred for a neurosurgical consultation and had previously sustained a work-related injury that resulted in herniation of a lumbar disc at one or more levels. The diagnosis in each case was confirmed by magnetic resonance imaging. The patient's chosen were scheduled for percutaneous discectomy. Introduction of the cannula for the purpose of performing percutaneous discectomy offered an opportunity to measure pressure changes in the disc prior to the operative procedure.

The patient was prepared and a cannula was inserted under local anesthesia into the nucleus pulposus of the L4-5 intervertebral disc, using anteroposterior and lateral fluoroscopy to position the end. With the cannula in place, the patient was moved to a VAX-D table. The VAX-D equipment* is routinely utilized in our non-surgical treatment program for patients suffering from low-back pain. The equipment consists of a split table design with a tensionometer mounted on the caudal, moveable section. The patient lies in a prone position and grasps handgrips to restrain movement of the upper body, which is supported on the fixed section of the table (Fig. I).



Fig. 1. Photograph illustrating the equipment and the position of the patient as the system is activated. The caudal end of the Table extends, applying tension to the pelvic belt. Upper body movement is restrained by having the patient grasp the handgrips. A graph of the tension applied is plotted by a chart recorder on the control console and the intradiscal pressure readings are entered on the same graph at the apex of each distraction curve.

The cannula was then connected to a pressure monitor using a disposable pressure transducer. The lines were filled with normal saline. The pelvic harness designed for this therapy was fastened around the pelvic girdle and connected to the tensionometer via straps attached to the harness. When the system was activated the caudal section supporting the lower body extended slowly, applying a distraction force via the pelvic harness connected to the tensionometer. The level of tension was preset by the operator on the control console and observed and plotted on a chart recorder. The movement of the table was stopped and held when the desired tension was reached. An average course of therapy consisted of 30-minute sessions on the table once a day for 10 to 15 days. During each session the patient undergoes alternating cycles of distraction and relaxation, the timing and periodicity having been programmed by the therapist.

In this study various distraction tensions, ranging from 50 to 100 lbs, were used for vertebral axial decompression therapy. The distraction tensions applied were monitored on a digital readout and recorded on a continuous graph tracing by a chart printer incorporated in the control console. The resulting changes in intradiscal pressure in the L4-5 nucleus pulposus were observed on a digital readout on the pressure monitor, and the readings were entered onto the chart recording at the point when the apex of distraction tension was achieved. The pressure readings were then applied to the negative-range calibrated curves prepared for each transducer to derive accurate intradiscal pressure readings.

The biological transducers employed in this study are primarily designed to measure pressure changes in the positive range. Following each procedure the pressure monitor and the disposable pressure transducer used for each patient were individually calibrated and a correction curve was plotted showing the transducer readings versus actual pressures, to correct for the nonlinearity of the instrumentation in the range of negative pressures

achieved. A pneumatic calibration analyzer with an accuracy of + 2% was used for this purpose.

Results

Intradiscal pressure measurements showed that distraction tension routinely applied by the VAX-D equipment reduced the intradiscal pressure significantly, to negative levels in the range of -100 to -160 mm Hg. The relationship between distraction tensions and intradiscal pressure changes for three patients is presented in Table 1. The extent of decompression (measured in mm Hg) shows an inverse relationship to the tension applied and may be expressed by a polynomial equation (Fig. 2).






FIG. 2. Graphs showing the intradiscal pressures recorded in the L4-5 nucleus pulposus of three patients (Case 3, upper; Case 4, center, and Case 5, lower) with a herniated disc at this level . Pressure is plotted against distraction tension consistent with the range of tension recommended as the therapeutic protocol for the equipment used in this study

Discussion

Intradiscal pressure changes were monitored in five patients. When the first two patients were tested, it was not recognized that biological transducers produce nonlinear measurements in the negative ranges at the levels achieved in this study. Since the disposable units had been discarded it was not possible to translate the findings accurately; however, the intradiscal pressures were observed to be significantly lowered. Also, the findings were consistent with the later three patients, for whom the transducers were retained and individually calibrated, permitting accurate interpretation of the results.

An interesting observation was that changes in intradiscal pressure appeared to be minimal until a threshold distraction tension was reached. When the threshold was exceeded the intradiscal pressure was observed to decrease dramatically, to levels in excess of 200 mm Hg below the positive pressure observed prior to the application of pelvic tension. As indicated in the curves plotted for intradiscal pressures versus distraction tension (Fig. 2), it appeared that the decrease in pressure tends to level off as the applied distraction tensions approached 100 lbs. The concept of a threshold distraction tension and the levels observed in these trials are consistent with radiographic studies of vertebral body separation reported in other publications²

The results indicate that it is possible to lower pressure in the nucleus pulposus of herniated lumbar discs to levels significantly below 0 mm Hg when distraction tension is applied according to the protocol described for vertebral axial decompression therapy. These findings may offer a plausible explanation for the mechanism of action for this therapeutic modality. Future research is warranted to study the decompression phenomenon achieved with this technology and its relationship to clinical outcome in patients with anatomical dysfunction of the lumbar spine. We are preparing a follow-up study on the clinical efficacy of this treatment modality.

Acknowledgments: The authors gratefully acknowledge the director and staff of the Department of Physical Therapy for their assistance in administering the vertebral axial decompression therapy.

Disclosure: The authors have no financial interest in either the equipment or the methodology advanced in this study.

References

1. Anderson GBJ, Schultz AB, Nachemson AL: Intervertebral disc pressures during traction. *Scand J Rehabil Suppl* 9:88-91, 1983
2. Colachis SC Jr, Strohm BR: Effects of intermittent traction on separation of lumbar vertebrae. *Arch Phys Med Rehabil* 50:251-258, 1969
3. Gupta RC, Ramarao SV: Epidurography in reduction of lumbar disc prolapse by traction. *Arch Phys Med Rehabil* 59:322-327, 1978
4. Hirsch C, Nachemson A: The reliability of lumbar disk surgery. *Clin Orthop* 29:189-195, 1963
5. Mathews JA: Dynamic discography: a study of lumbar traction. *Ann Phys Med* 9:275-279, 1968
6. Nachemson A: The load on lumbar disks in different positions of the body. *Clin Orthop* 45:107-122, 1966
7. Nachemson A: The lumbar spine-an orthopedic challenge. *Spine* 1:59, 1976
8. Nachemson A, Elfstrom G: Intravital dynamic pressure measurements in lumbar discs. A study of common movements, maneuvers and exercises. *Scand J Rehabil Med* 2:1, 1971
9. Nachemson AL: Disc pressure measurements. *Spine* 6:93-97, 1981
10. Pal B, Mangion P, Hossain MA, et al: A controlled trial of continuous lumbar traction in the treatment of back pain and sciatica. *Br J Rheumatol* 25:181-183 1986
11. Panjabi M, Brown M, Lindahl S. et al: Intrinsic disc pressure as a measure of integrity of the lumbar spine. *Spine* 13: 913-917, 1988
12. Spangfort EV: The lumbar disc herniation. A computer aided analysis of 2,504 operations. *Acta Orthop Scand Suppl* 142: 1-95, 1972
13. Tile M: The role of surgery in nerve root compression. *Spine* 9:57-64, 1984
14. Twomey LT: Sustained lumbar traction. An experimental study of long spine segments. *Spine* 10:146-149, 1985
15. Weber H: Lumbar disc herniation. A controlled prospective study with ten years of observation. *Spine* 8: 131 -140,1983